

PCMI: \mathbb{A}^1 -HOMOTOPY THEORY AND THE WEIL CONJECTURES PROBLEM SESSIONS

KIRSTEN WICKELGREN

CONTENTS

1. Problem Session for Lecture 1	1
2. Problem Session for Lecture 2	1
References	2

1. PROBLEM SESSION FOR LECTURE 1

Problem 1. Let X be a dualizable object of a symmetric monoidal category $(\mathcal{C}, \otimes, 1, \tau)$ with dual $\mathbb{D}X$. Show there is a natural bijection

$$\text{Map}(X \otimes Y, Z) = \text{Map}(Y, \mathbb{D}X \otimes Z)$$

Problem 2. Compute the categorical trace of a rational function on \mathbb{P}^1 , e.g. $z \mapsto z^3$, $z \mapsto z^2 + z + 1$, $z \mapsto \frac{z^5 + 1}{z^2 + 2z + 1}$

Problem 3. Compute the Zeta function of \mathbb{P}^n over a finite field.

Problem 4. Show that $\text{Th}(\mathbf{P}^1, \mathcal{O}(2)) \simeq \text{Th}(\mathbf{P}^1, \mathcal{O}(-2)) \simeq (\mathbf{P}^1)^{\wedge 2} \vee \mathbf{P}^1$.

Problem 5. [Sil09, V §2] Let E be an elliptic curve over a finite field and let F denote the relative Fröbenius. Show the Zeta function of E is of the form

$$\frac{(1 - \alpha T + qT^2)}{(1 - T)(1 - qT)}$$

as follows. Let $T_\ell E$ denote the Tate module. Let $\text{End}(E) \rightarrow \text{End}(T_\ell E)$ be denoted by $\psi \mapsto \psi_\ell$. One can use the Tate pairing to show that $\det \psi_\ell = \deg \psi$. $E(\mathbb{F}_{q^m}) = \deg(1 - F^m)$. Compute $\det(T - F_\ell^m)$ in terms of roots of $\det(T - F_\ell)$.

2. PROBLEM SESSION FOR LECTURE 2

Problem 6. Let u be a non-square in \mathbb{F}_{q^m} . Compute $\text{Tr}_{\mathbb{F}_{q^m}/\mathbb{F}_q} \langle u \rangle$ for $m = 1, 2, 3, \dots$

Problem 7. Compute the logarithmic \mathbb{A}^1 -zeta function of \mathbb{P}^n over a finite field.

Problem 8. Check that a symmetric monoidal functor (also called a \otimes -functor) $H : \mathcal{C} \rightarrow \mathcal{D}$ takes dualizable objects to dualizable objects and $\text{Tr } H = H \text{ Tr}$.

Date: July 14, 2024.

Problem 9. (See e.g. [Mil13, Lemma 27.5]) Let P be the characteristic polynomial of an endomorphism F of a finite dimensional vector space. Suppose $P(t) = \prod_r (1 - \alpha_r t)$. Show that $\text{Tr}(F^m) = \sum \alpha_i^m$ and

$$d \log P(t) = - \sum_m \text{Tr}(F^m) t^{m-1},$$

where $d \log P(t) = P'(t)/P(t)$.

Problem 10. [Mor12, Lemma 3.5] We've asserted that $\text{GW}(k) \cong K_\delta^{\text{MW}}(k)$. This exercise is to help get comfortable with that. We define $K_*^{\text{MW}}(k)$ as the graded associative ring generated by symbols $[a]$ for each a in k^* of degree 1 and a symbol η of degree -1 subject to the relations

- (1) $[a][1 - a] = 0$
- (2) $[ab] = [a] + [b] + \eta[a][b]$
- (3) $\eta[a] = [a]\eta$
- (4) $h\eta = 0$ where $h = 1 + 1 + \eta[-1]$.

Define $\langle a \rangle$ in $K_\delta^{\text{MW}}(k)$ by $\langle a \rangle = 1 + \eta[a]$. Show that $[ab] = [a] + \langle a \rangle[b] = [a]\langle b \rangle + [b]$, and that $\langle ba \rangle = \langle a \rangle \langle b \rangle$.

REFERENCES

- [Mil13] J.S. Milne, *Lectures on etale cohomology* (2013). Available at <https://www.jmilne.org/math/CourseNotes/LEC.pdf>. $\uparrow 9$
- [Mor12] Fabien Morel, *\mathbb{A}^1 -algebraic topology over a field*, Lecture Notes in Mathematics, vol. 2052, Springer, Heidelberg, 2012. MR2934577 $\uparrow 10$
- [Sil09] Joseph H. Silverman, *The arithmetic of elliptic curves*, Second, Graduate Texts in Mathematics, vol. 106, Springer, Dordrecht, 2009. MR2514094 $\uparrow 5$

K. WICKELGREN, DEPARTMENT OF MATHEMATICS, DUKE UNIVERSITY, 120 SCIENCE DR, DURHAM NC

Email address: kirsten.wickelgren@duke.edu

URL: <https://services.math.duke.edu/~kgw/>