

Constructions of Expanders Using Group Theory

Martin Kassabov,
Cornell University and IAS

November 2009

Overview

Expander Graphs

Kazhdan Constants

Computation of Kazhdan Constants

Bounded Generation Property

Kazhdan Constants for $SL_n(\mathbb{F}_p)$

Using Geometry to (dis-)Prove Property T

Subspace Arrangements

Easy Example

Applications

Expander Graphs

Kazhdan Constants

Computation of Kazhdan Constants

Bounded Generation Property

Kazhdan Constants for $SL_n(\mathbb{F}_p)$

Using Geometry to (dis-)Prove Property T

Subspace Arrangements

Easy Example

Applications

Expander Graphs

- Expanders
- Equivalent definitions
- Cayley Graphs

Kazhdan Constants

Computation of Kazhdan Constants

Bounded Generation Property

Kazhdan Constants for $SL_n(\mathbb{F}_p)$

Using Geometry to (dis-)Prove Property T

Subspace Arrangements

Easy Example

Applications

Expander Graphs

Expanders

Expander Graphs

● **Expanders**

● Equivalent definitions

● Cayley Graphs

Kazhdan Constants

Computation of Kazhdan Constants

Bounded Generation Property

Kazhdan Constants for $SL_n(\mathbb{F}_p)$

Using Geometry to (dis-)Prove Property T

Subspace Arrangements

Easy Example

Applications

Well connected graphs of small degree are called expanders.

Definition *A graph Γ is called an ε -expander if for any set of vertices A , such that $|A| \leq |\Gamma|/2$ we have $|\partial A| \geq \varepsilon|A|$.*

The maximal ε with this property is called the expanding constant of Γ .

A family of graphs (of bounded degree) is called an expander family if their expanding constants are uniformly bounded.

Equivalent definitions of Expanders

Expander Graphs

- Expanders
- **Equivalent definitions**
- Cayley Graphs

Kazhdan Constants

Computation of Kazhdan Constants

Bounded Generation Property

Kazhdan Constants for $SL_n(\mathbb{F}_p)$

Using Geometry to (dis-)Prove Property T

Subspace Arrangements

Easy Example

Applications

Theorem *A family of k -regular graphs Γ_i are expanders if any of the following holds:*

- *the expanding constants of Γ_i are bounded from 0,*
- *the Cheeger constants of Γ_i are bounded from 0,*
- *the spectral gaps of the Laplacians ($\mathbb{1}_d$ -normalized adjacency matrices) are bounded from 0.*

Any of these conditions implies that the random walks on Γ_i mix in $O(\log |\Gamma_i|)$ steps.

If one allows k to increase then these conditions are NOT equivalent, thus there are several different definitions of unbounded degree expanders.

Cayley Graphs as Expanders

Expander Graphs

- Expanders
- Equivalent definitions
- **Cayley Graphs**

Kazhdan Constants

Computation of Kazhdan Constants

Bounded Generation Property

Kazhdan Constants for $SL_n(\mathbb{F}_p)$

Using Geometry to (dis-)Prove Property T

Subspace Arrangements

Easy Example

Applications

In the case of Cayley graphs of finite groups, there is another condition equivalent to “expansion”, which is related to representation theory, more precisely to the “Kazhdan constants”.

Problem *Let be $\{G_i\}$ a family of finite groups. Is it possible to find generating sets S_i , which make the Cayley graphs $\mathcal{C}(G_i, S_i)$ expanders?*

- there are many cases where such generating sets are known to exist, e.g., for quotients of a group with a variant of property T;
- and only a few where it can be proven that such generating sets do not exist, e.g., solvable groups of a fixed class.

Expander Graphs

Kazhdan Constants

- Unitary Representations
- Kazhdan Property T
- Examples
- Several Kazhdan Constants

Computation of Kazhdan Constants

Bounded Generation Property

Kazhdan Constants for $SL_n(\mathbb{F}_p)$

Using Geometry to (dis-)Prove Property T

Subspace Arrangements

Easy Example

Applications

Kazhdan Constants

Unitary Representations

Expander Graphs

Kazhdan Constants

• Unitary Representations

• Kazhdan Property T

• Examples

• Several Kazhdan Constants

Computation of Kazhdan Constants

Bounded Generation Property

Kazhdan Constants for $SL_n(\mathbb{F}_p)$

Using Geometry to (dis-)Prove Property T

Subspace Arrangements

Easy Example

Applications

Let \mathcal{H} be a Hilbert space.

A representation a group of G into \mathcal{H} is called unitary if

$$\|g(v)\| = \|v\|$$

for any $v \in \mathcal{H}$ and $g \in G$.

Definition A unit vector $v \in \mathcal{H}$ is called ε -almost invariant under the set S if

$$\|g(v) - v\| \leq \varepsilon \text{ for any } g \in S.$$

One easy way to produce almost invariant vectors is to take perturbations of invariant vectors.

Kazhdan Property T

Expander Graphs

Kazhdan Constants

• Unitary

Representations

• Kazhdan Property T

• Examples

• Several Kazhdan

Constants

Computation of Kazhdan

Constants

Bounded Generation

Property

Kazhdan Constants for

$SL_n(\mathbb{F}_p)$

Using Geometry to

(dis-)Prove Property T

Subspace Arrangements

Easy Example

Applications

Definition Let G be a group, generated by a finite set S .

The maximal ε such that the existence of an ε -almost invariant vector implies the existence of an invariant vector is called Kazhdan constant of G with respect to S and is denoted by $\mathcal{K}(G; S)$.

The group G has property T if the Kazhdan constant $\mathcal{K}(G; S)$ is positive.

If H is a normal subgroup of G , we can also define about *relative Kazhdan constants*:

Definition The maximal ε such that the existence of an ε -almost invariant vector implies the existence of an H invariant vector is called relative Kazhdan constant of G with respect to S relative to H and is denoted by $\mathcal{K}(G, H; S)$.

Examples of Groups with T

Expander Graphs

Kazhdan Constants

- Unitary Representations
- Kazhdan Property T
- **Examples**
- Several Kazhdan Constants

Computation of Kazhdan Constants

Bounded Generation Property

Kazhdan Constants for $SL_n(\mathbb{F}_p)$

Using Geometry to (dis-)Prove Property T

Subspace Arrangements

Easy Example

Applications

- All finite groups
- Lattices in high rank Lie groups, e.g., $SL_n(\mathbb{Z})$
- Random group, with sufficiently many relations

Infinite groups with property T can be used to construct expander graphs.

Also, families of finite groups with a generating sets of fixed size lead to expanders, provided that their Kazhdan constants are uniformly bounded.

Moreover, estimates for the Kazhdan constant give to estimates for the mixing time for some random walks.

Several Kazhdan Constants

Expander Graphs

Kazhdan Constants

- Unitary Representations
- Kazhdan Property T
- Examples
- **Several Kazhdan Constants**

Computation of Kazhdan Constants

Bounded Generation Property

Kazhdan Constants for $SL_n(\mathbb{F}_p)$

Using Geometry to (dis-)Prove Property T

Subspace Arrangements

Easy Example

Applications

Let G be a group, generated by a finite set S .

The Kazhdan constant of G is

$$\mathcal{K}(G; S) = \inf_{\mathcal{H}} \inf_{v \in \mathcal{H}} \max_{s \in S} \frac{\|\rho(s)v - v\|}{\|v\|}$$

where \mathcal{H} is a representation of G without invariant vectors

Similarly we can define

$$\mathcal{K}_{av}^2(G; S) = \inf_{\mathcal{H}} \inf_{v \in \mathcal{H}} \frac{1}{|S|} \sum_{s \in S} \frac{\|\rho(s)v - v\|^2}{\|v\|^2}$$

$\mathcal{K}_{av}^2(G; S)$ is equal to twice the spectral gap of the Laplacian and the inf is achieved at some irreducible representation.

Expander Graphs

Kazhdan Constants

Computation of Kazhdan Constants

- Trivial bounds
- Dependence on the generating set
- Some Relative Kazhdan Constants

Bounded Generation Property

Kazhdan Constants for $SL_n(\mathbb{F}_p)$

Using Geometry to (dis-)Prove Property T

Subspace Arrangements

Easy Example

Applications

Computation of Kazhdan Constants

Trivial bounds

Expander Graphs

Kazhdan Constants

Computation of Kazhdan Constants

● **Trivial bounds**

● Dependence on the generating set

● Some Relative Kazhdan Constants

Bounded Generation Property

Kazhdan Constants for $SL_n(\mathbb{F}_p)$

Using Geometry to (dis-)Prove Property T

Subspace Arrangements

Easy Example

Applications

It is very easy to see that:

$$2 \geq \mathcal{K}(G; G) > \sqrt{2},$$

and

$$\mathcal{K}_{av}^2(G; G) = 2.$$

These inequalities hold, because if unit vector v is moved by less than $\sqrt{2}$ by any element of the group G then the whole orbit of v lies in some half-space.

Thus, the center of mass of the orbit is a non-zero invariant vector in the representation.

Equivalently, we use that the normalized adjacency matrix of a complete graph has a spectral gap equal to 1.

Dependance on the generating set I

Let G be a group and let S be a set (containing the identity) then

$$\mathcal{K}(G; S) \geq \frac{1}{k} \mathcal{K}(G; S^k),$$

where S^k denotes all group elements which can be written as a product of k elements from S .

This inequality holds, because any ϵ -almost invariant vector w.r.t. S is also $k\epsilon$ -almost invariant w.r.t. to S^k .

Expander Graphs

Kazhdan Constants

Computation of Kazhdan Constants

- Trivial bounds
- **Dependance on the generating set**

• Some Relative Kazhdan Constants

Bounded Generation Property

Kazhdan Constants for $SL_n(\mathbb{F}_p)$

Using Geometry to (dis-)Prove Property T

Subspace Arrangements

Easy Example

Applications

Dependance on the generating set II

Expander Graphs

Kazhdan Constants

Computation of Kazhdan Constants

- Trivial bounds
- **Dependance on the generating set**
- Some Relative Kazhdan Constants

Bounded Generation Property

Kazhdan Constants for $SL_n(\mathbb{F}_p)$

Using Geometry to (dis-)Prove Property T

Subspace Arrangements

Easy Example

Applications

If H is a normal subgroup of G then

$$\mathcal{K}(G; S) \geq \frac{1}{2} \mathcal{K}(G; H \cup S) \mathcal{K}(G, H; S),$$

where $\mathcal{K}(G, H; S)$ is the relative Kazhdan constant.

This inequality holds, because any ϵ -almost invariant vector w.r.t. S is also $2\mathcal{K}(G, H; S)^{-1}\epsilon$ -almost invariant w.r.t. to H .

Lemma *If $H_i \triangleleft N_i$ are subgroups of G such that*

$$\mathcal{K}(N_i, H_i; S_i) \geq L.$$

Then

$$\mathcal{K}(G; \cup S_i) \geq \frac{1}{2} L \times \mathcal{K}(G; \cup H_i).$$

Some Relative Kazhdan Constants

Expander Graphs

Kazhdan Constants

Computation of Kazhdan Constants

- Trivial bounds
- Dependence on the generating set

• **Some Relative Kazhdan Constants**

Bounded Generation Property

Kazhdan Constants for $SL_n(\mathbb{F}_p)$

Using Geometry to (dis-)Prove Property T

Subspace Arrangements

Easy Example

Applications

Theorem (Burger, Shalom, K.)

Let R be an associative ring generated by $\alpha_i, i = 1, \dots, k$.

Consider the subgroup $GL_2(R) \rtimes R^2 \subset GL_3(R)$.

$$\begin{pmatrix} * & * & * \\ * & * & * \\ 0 & 0 & 1 \end{pmatrix}$$

Then

$$\mathcal{K}(GL_2(R) \rtimes R^2, R^2; F) \geq \frac{1}{5(\sqrt{k} + 3)},$$

where F is the set consisting of $4(k + 1)$ elementary matrices with ± 1 or $\pm \alpha_i$ off the diagonal and the two standard generators of R^2 .

The proof uses almost invariant measures on the unitary dual of R^2 considered as additive group.

Expander Graphs

Kazhdan Constants

Computation of Kazhdan
Constants

**Bounded Generation
Property**

- Bounded Generation
- Example: $SL_n(\mathbb{Z})$
- Uniform bounded elementary generation

Kazhdan Constants for
 $SL_n(\mathbb{F}_p)$

Using Geometry to
(dis-)Prove Property T

Subspace Arrangements

Easy Example

Applications

Bounded Generation Property

Bounded Generation

Expander Graphs

Kazhdan Constants

Computation of Kazhdan Constants

Bounded Generation Property

● **Bounded Generation**

- Example: $SL_n(\mathbb{Z})$
- Uniform bounded elementary generation

Kazhdan Constants for $SL_n(\mathbb{F}_p)$

Using Geometry to (dis-)Prove Property T

Subspace Arrangements

Easy Example

Applications

Definition A group G is said to be boundedly generated by the ordered multi-set $S = \{s_1, s_2, \dots, s_k\}$ if any element $g \in G$ can be written as a product

$$g = s_1^{a_1} s_2^{a_2} \dots s_k^{a_k}$$

for some integers a_i .

This is a very strong condition and implies strong restrictions on the group G .

Example: $SL_n(\mathbb{Z})$

Expander Graphs

Kazhdan Constants

Computation of Kazhdan Constants

Bounded Generation Property

- Bounded Generation
- **Example: $SL_n(\mathbb{Z})$**
- Uniform bounded elementary generation

Kazhdan Constants for $SL_n(\mathbb{F}_p)$

Using Geometry to (dis-)Prove Property T

Subspace Arrangements

Easy Example

Applications

Theorem (Carter-Keller) *Any matrix in $SL_n(\mathbb{Z})$ can be written as a product of less than $2n^2 + 100$ elementary matrices.*

The proof uses reduction from $SL_n(\mathbb{Z})$ to $SL_{n-1}(\mathbb{Z})$ for $n \geq 3$, which only uses existence of primes in arithmetic progression.

The main step is to show that any element in $SL_2(\mathbb{Z}) \subset SL_3(\mathbb{Z})$ can be written as a product of 100 elementary matrices.

This step uses K-theory and playing with Mennicke symbols.

If we are only interested in $SL_n(\mathbb{F}_p)$ then the proof becomes much easier and uses only Gauss elimination.

Uniform bounded elementary generation

Expander Graphs

Kazhdan Constants

Computation of Kazhdan Constants

Bounded Generation Property

- Bounded Generation
- Example: $SL_n(\mathbb{Z})$
- **Uniform bounded elementary generation**

Kazhdan Constants for $SL_n(\mathbb{F}_p)$

Using Geometry to (dis-)Prove Property T

Subspace Arrangements

Easy Example

Applications

For groups like $SL_n(R)$ it is better to consider the following property

Definition A group G called is boundedly generated by the subgroups H_i , if there exists N such that any $g \in G$ can be written as

$$g = g_1 g_2 \dots g_N, \quad \text{where } g_i \in \bigcup H_i.$$

Theorem Let R be a “nice” finite associative ring with 1, then any element in $EL_3(R)$ can be written as a product of 50 elementary matrices.

A “nice” ring is any product of matrix rings over fields, or commutative rings.

It is unknown if similar result holds for $SL_n(\mathbb{Z}[x])$ for a big n .
It is false for $SL_n(\mathbb{C}[x])$.

Expander Graphs

Kazhdan Constants

Computation of Kazhdan Constants

Bounded Generation Property

Kazhdan Constants for $SL_n(\mathbb{F}_p)$

- $SL_n(\mathbb{F}_p)$ for fixed n
- $SL_{3n}(\mathbb{F}_p)$ for all n and p

Using Geometry to (dis-)Prove Property T

Subspace Arrangements

Easy Example

Applications

Kazhdan Constants for $SL_n(\mathbb{F}_p)$

$SL_n(\mathbb{F}_p)$ for fixed $n \geq 3$

Expander Graphs

Kazhdan Constants

Computation of Kazhdan Constants

Bounded Generation Property

Kazhdan Constants for $SL_n(\mathbb{F}_p)$

• $SL_n(\mathbb{F}_p)$ for fixed n

• $SL_{3n}(\mathbb{F}_p)$ for all n and p

Using Geometry to (dis-)Prove Property T

Subspace Arrangements

Easy Example

Applications

The homomorphisms from $SL_2(\mathbb{Z}) \times \mathbb{Z}^2$ to $SL_n(\mathbb{F}_p)$, and estimates for the relative Kazhdan constant of $SL_2(\mathbb{Z}) \times \mathbb{Z}^2$ imply that

$$\mathcal{K}(SL_n(\mathbb{F}_p); E_n(\pm 1)) \geq \frac{1}{20} \mathcal{K}(SL_n(\mathbb{F}_p); E_n),$$

where $E_n(\pm 1)$ is the set of elementary matrices with ± 1 off the diagonal and E_n is the set of all el. matrices.

By bounded generation $E_n^{2n^2} = SL_n$, therefore

$$\mathcal{K}(SL_n(\mathbb{F}_p); E_n) \geq \frac{1}{2n^2} \mathcal{K}(SL_n(\mathbb{F}_p); SL_n(\mathbb{F}_p)) \geq \frac{1}{2n^2}$$

Theorem (Shalom)

$$\mathcal{K}(SL_n(\mathbb{F}_p); E_n(\pm 1)) > \frac{1}{40n^2}.$$

$SL_{3n}(\mathbb{F}_p)$ for a all n and p

Expander Graphs

Kazhdan Constants

Computation of Kazhdan Constants

Bounded Generation Property

Kazhdan Constants for $SL_n(\mathbb{F}_p)$

• $SL_n(\mathbb{F}_p)$ for fixed n

• $SL_{3n}(\mathbb{F}_p)$ for a all n and p

Using Geometry to (dis-)Prove Property T

Subspace Arrangements

Easy Example

Applications

Block decomposition implies $SL_{3n}(\mathbb{F}_p) \simeq EL_3(\text{Mat}_n(\mathbb{F}_p))$.

Since the ring $\text{Mat}_n(\mathbb{F}_p)$ is 2-generated, and $EL_3(\text{Mat}_n(\mathbb{F}_p))$ has uniform bounded generation property, we have:

$$\mathcal{K}(SL_{3n}(\mathbb{F}_p); E_3(\pm 1, A, B)) \geq \frac{1}{20} \mathcal{K}(SL_{3n}(\mathbb{F}_p); E_3),$$

$$\mathcal{K}(SL_{3n}(\mathbb{F}_p); E_3) \geq \frac{1}{40} \mathcal{K}(SL_{3n}(\mathbb{F}_p); SL_{3n}(\mathbb{F}_p)) \geq \frac{1}{40}$$

where $E_3(\pm 1, A, B)$ is the set of block elementary matrices with generators of $\text{Mat}_n(\mathbb{F}_p)$ off the diagonal and E_3 is the set of all block el. matrices.

Theorem (K) *There resists a generating set S of $SL_{3n}(\mathbb{F}_p)$ with $|S| < 20$ and*

$$\mathcal{K}(SL_{3n}(\mathbb{F}_p); S) \geq \frac{1}{1000}.$$

Expander Graphs

Kazhdan Constants

Computation of Kazhdan Constants

Bounded Generation Property

Kazhdan Constants for $SL_n(\mathbb{F}_p)$

Using Geometry to (dis-)Prove Property T

- Example D_∞
- Groups generated by involutions
- Angle between two subspaces
- Angle between many subspaces
- Bounding the angle between subspaces
- Groups generated by 3 involutions
- 3 Subspaces

Subspace Arrangements

Easy Example

Applications

Using Geometry to (dis-)Prove Property T

Example D_∞

Expander Graphs

Kazhdan Constants

Computation of Kazhdan Constants

Bounded Generation Property

Kazhdan Constants for $SL_n(\mathbb{F}_p)$

Using Geometry to (dis-)Prove Property T

● **Example D_∞**

- Groups generated by involutions
- Angle between two subspaces
- Angle between many subspaces
- Bounding the angle between subspaces
- Groups generated by 3 involutions
- 3 Subspaces

Subspace Arrangements

Easy Example

Applications

The infinite dihedral group


$$D_\infty = \langle a, b \mid a^2 = b^2 = 1 \rangle$$

does not have property T.

For any φ there is a representation of D_∞ on \mathbb{R}^2 where a, b act as reflections across two lines at angle 2φ .

This representation does not have invariant vectors, but it has a unit vector which is $2 \sin \varphi$ invariant.

This example shows that we need some bound for the angle between the subspaces \mathcal{H}^{G_1} and \mathcal{H}^{G_2} in order to prove property T for $G = \langle G_1, G_2 \rangle$.



Groups generated by involutions

Expander Graphs

Kazhdan Constants

Computation of Kazhdan Constants

Bounded Generation Property

Kazhdan Constants for $SL_n(\mathbb{F}_p)$

Using Geometry to (dis-)Prove Property T

- Example D_∞
- **Groups generated by involutions**
- Angle between two subspaces
- Angle between many subspaces
- Bounding the angle between subspaces
- Groups generated by 3 involutions
- 3 Subspaces

Subspace Arrangements

Easy Example

Applications

Assume that the generating set S consist of involutions.

Let \mathcal{H}_s denotes the fixed subspace of $s \in S$.

Notice that:

- \mathcal{H} has no G invariant vectors $\Leftrightarrow \bigcap \mathcal{H}_s = \{0\}$
- $\rho(s)v - v = 2d(v, \mathcal{H}_s)$

Thus, the group G has property T if for any representation of G on \mathcal{H} , any vector which is close to all subspaces \mathcal{H}_s is not too far from the intersection $\bigcap \mathcal{H}_s$.

Angle between two subspaces

Expander Graphs

Kazhdan Constants

Computation of Kazhdan Constants

Bounded Generation Property

Kazhdan Constants for $SL_n(\mathbb{F}_p)$

Using Geometry to (dis-)Prove Property T

- Example D_∞
- Groups generated by involutions
- **Angle between two subspaces**
- Angle between many subspaces
- Bounding the angle between subspaces
- Groups generated by 3 involutions
- 3 Subspaces

Subspace Arrangements

Easy Example

Applications

Definition Let \mathcal{H}_1 and \mathcal{H}_2 be two subspaces of a Hilbert space \mathcal{H} . The angle $\sphericalangle(\mathcal{H}_1, \mathcal{H}_2)$ is the smallest angle between vectors v_1 and v_2 , such that $v_i \in \mathcal{H}_i$ and $v_i \perp \mathcal{H}_1 \cap \mathcal{H}_2$.

Notice that the operator norm of the addition

$$\text{sum} : \mathcal{H}_1 / (\mathcal{H}_1 \cap \mathcal{H}_2) \oplus \mathcal{H}_2 / (\mathcal{H}_1 \cap \mathcal{H}_2) \rightarrow \mathcal{H} / (\mathcal{H}_1 \cap \mathcal{H}_2)$$

is less than $\sqrt{1 + \cos \sphericalangle(\mathcal{H}_1, \mathcal{H}_2)}$.

Equivalently, we can define the angle using the spectral gap near $\sqrt{2}$ of the addition

$$\text{sum} : \mathcal{H}_1 \oplus \mathcal{H}_2 \rightarrow \mathcal{H}$$

Notice that the angle between two subspaces is 'not defined' if one is subspace of the other. In this case we say that the angle is $\pi/2$ and the subspaces are perpendicular.

Angle between many subspaces

Expander Graphs

Kazhdan Constants

Computation of Kazhdan Constants

Bounded Generation Property

Kazhdan Constants for $SL_n(\mathbb{F}_p)$

Using Geometry to (dis-)Prove Property T

- Example D_∞
- Groups generated by involutions
- Angle between two subspaces
- **Angle between many subspaces**
- Bounding the angle between subspaces
- Groups generated by 3 involutions
- 3 Subspaces

Subspace Arrangements

Easy Example

Applications

Definition Let $\mathcal{H}_i, i = 1, \dots, n$ be subspaces of a Hilbert space \mathcal{H} . We say that the angle $\angle(\mathcal{H}_i)$ between them is ϕ if

$$\cos \phi = \frac{1}{n-1} \left(\sup \frac{\|\sum v_i\|^2}{\sum \|v_i\|^2} - 1 \right)$$

where sup is taken over $v_i \in \mathcal{H}_i$ and $v_j \perp \bigcap_i \mathcal{H}_i$.

Equivalently the norm of the addition

$$\text{sum} : \mathcal{H}_1 / \bigcap \mathcal{H}_i \oplus \dots \mathcal{H}_n / \bigcap \mathcal{H}_i \rightarrow \mathcal{H} / \bigcap \mathcal{H}_i$$

is $\sqrt{1 + (n-1) \cos \phi}$.

Notice that the angle between any collection of subspaces in a finite dimensional spaces is positive, but this is not true in general.

Also any lower bound for $\angle(\mathcal{H}_i)$ gives a lower bound of maximal distance from a unit vector v to the subspaces \mathcal{H}_i .

Bounding the angle between subspaces

Expander Graphs

Kazhdan Constants

Computation of Kazhdan Constants

Bounded Generation Property

Kazhdan Constants for $SL_n(\mathbb{F}_p)$

Using Geometry to (dis-)Prove Property T

- Example D_∞
- Groups generated by involutions
- Angle between two subspaces
- Angle between many subspaces
- **Bounding the angle between subspaces**
- Groups generated by 3 involutions
- 3 Subspaces

Subspace Arrangements

Easy Example

Applications

Lemma *If the involutions s and t generate a dihedral group of order $2k$ then the angle $\angle(\mathcal{H}_s, \mathcal{H}_t) \geq \pi/k$, where \mathcal{H}_s and \mathcal{H}_t are the fixed points of s and t in any unitary representation of G .*

The proof uses the representation theory of the dihedral group D_k .

Similar result hold if s and t are elements of order p and $\langle s, t \rangle$ is the Heisenberg group mod p . In this case we have

$$\cos \angle(\mathcal{H}_s, \mathcal{H}_t) \leq \frac{1}{\sqrt{p}}.$$

Again it suffices to verify this bound for any irreducible representation of the Heisenberg group.

Groups generated by 3 involutions

Let G be a group generated by 3 involutions, i.e., G is a Coxeter group

$$G = \langle s_1, s_2, s_3 \mid s_i^2 = 1, (s_i s_j)^{m_{ij}} = 1 \rangle$$

Question *When does G has property T?*

Using the classification of 3 generated Coxeter groups one can see that the following are equivalent:

- G has property T
- G is finite
- $\frac{1}{m_{12}} + \frac{1}{m_{23}} + \frac{1}{m_{31}} > 1$

Expander Graphs

Kazhdan Constants

Computation of Kazhdan Constants

Bounded Generation Property

Kazhdan Constants for $SL_n(\mathbb{F}_p)$

Using Geometry to (dis-)Prove Property T

- Example D_∞
- Groups generated by involutions
- Angle between two subspaces
- Angle between many subspaces
- Bounding the angle between subspaces
- **Groups generated by 3 involutions**
- 3 Subspaces

Subspace Arrangements

Easy Example

Applications

3 Subspaces

Expander Graphs

Kazhdan Constants

Computation of Kazhdan Constants

Bounded Generation Property

Kazhdan Constants for $SL_n(\mathbb{F}_p)$

Using Geometry to (dis-)Prove Property T

- Example D_∞
- Groups generated by involutions
- Angle between two subspaces
- Angle between many subspaces
- Bounding the angle between subspaces
- Groups generated by 3 involutions

• 3 Subspaces

Subspace Arrangements

Easy Example

Applications

Let \mathcal{H}_i be 3 subspaces in a Hilbert space \mathcal{H} , with trivial intersection.

Lemma: Let α_{ij} denote the angle between \mathcal{H}_i and \mathcal{H}_j . If $\sum \alpha_{ij} > \pi$ than any vector v which is closed to all \mathcal{H}_i , is short, i.e.,


$$\|v\|^2 \leq C \sum d(v, \mathcal{H}_i)^2,$$

where the constant C depends only in α_{ij} .

Notice that the condition

$$\sum \alpha_{ij} > \pi$$

is equivalent to the existence of spherical triangle with angles α_{ij} .



Expander Graphs

Kazhdan Constants

Computation of Kazhdan Constants

Bounded Generation Property

Kazhdan Constants for $SL_n(\mathbb{F}_p)$

Using Geometry to (dis-)Prove Property T

Subspace Arrangements

- Subspace Arrangements
- 3 Subspaces
- Relationship with groups
- Easy Observation

Easy Example

Applications

Subspace Arrangements

Subspace Arrangements

Expander Graphs

Kazhdan Constants

Computation of Kazhdan Constants

Bounded Generation Property

Kazhdan Constants for $SL_n(\mathbb{F}_p)$

Using Geometry to (dis-)Prove Property T

Subspace Arrangements

● Subspace Arrangements

● 3 Subspaces

● Relationship with groups

● Easy Observation

Easy Example

Applications

Definition A subspace arrangement is the collection of all possible n -tuples of subspaces $\{\mathcal{H}_i\}_{i \in I}$ in some Hilbert space \mathcal{H} , satisfying a fixed set of conditions of the following type.

- for some subsets $J \subset I$ there exists α_J such that

$$\angle(\mathcal{H}_j)_{j \in J} \geq \alpha_J$$

- for some subsets $J \subset I$ and some $i \in I$ we have $\mathcal{H}_i \supset \bigcap_{j \in J} \mathcal{H}_j$

A collection is called “good”, if there is a strictly positive lower bound for

$$\angle(\mathcal{H}_i)_{i \in I},$$

which is valid for all n -tuples of subspaces which satisfy the conditions listed above.

3 Subspaces

Expander Graphs

Kazhdan Constants

Computation of Kazhdan Constants

Bounded Generation Property

Kazhdan Constants for $SL_n(\mathbb{F}_p)$

Using Geometry to (dis-)Prove Property T

Subspace Arrangements

• Subspace Arrangements

• **3 Subspaces**

• Relationship with groups

• Easy Observation

Easy Example

Applications

The previous lemma is equivalent to saying that any 3 subspaces such that

- $\angle(\mathcal{H}_1, \mathcal{H}_2) \geq \alpha_{12}$
- $\angle(\mathcal{H}_2, \mathcal{H}_3) \geq \alpha_{23}$
- $\angle(\mathcal{H}_3, \mathcal{H}_1) \geq \alpha_{31}$

satisfy $\angle(\mathcal{H}_1, \mathcal{H}_2, \mathcal{H}_3) \geq \alpha_0 > 0$ if

$$\alpha_{12} + \alpha_{23} + \alpha_{31} > \pi.$$

In other words the arrangement of 3 subspaces \mathcal{H}_i , satisfying the above conditions is “good”.

Relationship with groups

Expander Graphs

Kazhdan Constants

Computation of Kazhdan Constants

Bounded Generation Property

Kazhdan Constants for $SL_n(\mathbb{F}_p)$

Using Geometry to (dis-)Prove Property T

Subspace Arrangements

● Subspace Arrangements

● 3 Subspaces

● Relationship with groups

● Easy Observation

Easy Example

Applications

Our motivation for studying such arrangements of subspaces comes from representation theory.

Let \mathcal{H} be a representation of a group G and let G_i be finite subgroups of G . Denote $\mathcal{H}_i = \mathcal{H}^{G_i}$ the set of vectors fixed by each subgroup.

The two type of restrictions on the subspaces are related to properties of the group G (and the collections of subgroups G_i):

- representation theory of the group $\langle G_{i_1}, \dots, G_{i_k} \rangle$ yields lower bounds for the angle $\angle(\mathcal{H}_{i_1}, \dots, \mathcal{H}_{i_k})$,
- inclusions $G_j \subset \langle G_{i_1}, \dots, G_{i_k} \rangle$ correspond $\mathcal{H}_j \supset \bigcap_k \mathcal{H}_{i_k}$.

Easy Observation

Expander Graphs

Kazhdan Constants

Computation of Kazhdan Constants

Bounded Generation Property

Kazhdan Constants for $SL_n(\mathbb{F}_p)$

Using Geometry to (dis-)Prove Property T

Subspace Arrangements

- Subspace Arrangements
- 3 Subspaces
- Relationship with groups
- **Easy Observation**

Easy Example

Applications

Theorem *Let G be a group generated by n finite subgroups G_i . Suppose that the inclusions between G_J for $J \subset \{1, \dots, n\}$ and the representation theory of these groups define a "good" arrangement of subspaces. Then G has property T.*

Expander Graphs

Kazhdan Constants

Computation of Kazhdan Constants

Bounded Generation Property

Kazhdan Constants for $SL_n(\mathbb{F}_p)$

Using Geometry to (dis-)Prove Property T

Subspace Arrangements

Easy Example

- An Example
- Main result
- Equivalent Statement
- “Proof”
- Application to property T

Applications

Easy Example

An Example

Expander Graphs

Kazhdan Constants

Computation of Kazhdan Constants

Bounded Generation Property

Kazhdan Constants for $SL_n(\mathbb{F}_p)$

Using Geometry to (dis-)Prove Property T

Subspace Arrangements

Easy Example

- **An Example**
- Main result
- Equivalent Statement
- “Proof”
- Application to property T

Applications

Suppose that a collection of subspaces is given only by conditions of the first type, in the simplest case we have bounds for the angles α_{ij} between any pair of \mathcal{H}_i -es.

If we find sufficient condition for having a good collection we can deduce property T for some groups.

Previous results: $(\varepsilon_{i,j} = \cos \alpha_{i,j})$

- Dymara-Januszkiewicz proved that the collection is “good” if $\varepsilon_{ij} < 2^{-12n}$ for all i, j
- Ershov-Jaikin improved the bound to $\varepsilon_{ij} < \frac{1}{n-1}$ for all i, j
- They also showed that the collection is good if $n = 3$ and some inequality is satisfied.

Main result

Expander Graphs

Kazhdan Constants

Computation of Kazhdan Constants

Bounded Generation Property

Kazhdan Constants for $SL_n(\mathbb{F}_p)$

Using Geometry to (dis-)Prove Property T

Subspace Arrangements

Easy Example

- An Example
- **Main result**
- Equivalent Statement
- “Proof”
- Application to property T

Applications

Theorem (K.) Let \mathcal{H}_i are subspaces of a Hilbert space \mathcal{H} . Denote $\varepsilon_{i,j} = \cos(\angle(\mathcal{H}_i, \mathcal{H}_j))$. Suppose that the matrix

$$A = \begin{pmatrix} 1 & -\varepsilon_{1,2} & -\varepsilon_{1,3} & \dots & -\varepsilon_{1,n} \\ -\varepsilon_{2,1} & 1 & -\varepsilon_{2,2} & \dots & -\varepsilon_{2,n} \\ -\varepsilon_{3,1} & -\varepsilon_{3,2} & 1 & \dots & -\varepsilon_{3,n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ -\varepsilon_{n,1} & -\varepsilon_{n,2} & -\varepsilon_{n,3} & \dots & 1 \end{pmatrix}$$

is positive definite, then there is lower bound for $\angle(\mathcal{H}_i)$, which depends only the matrix A .

Equivalent Statement

Expander Graphs

Kazhdan Constants

Computation of Kazhdan Constants

Bounded Generation Property

Kazhdan Constants for $SL_n(\mathbb{F}_p)$

Using Geometry to (dis-)Prove Property T

Subspace Arrangements

Easy Example

- An Example
- Main result
- **Equivalent Statement**
- “Proof”
- Application to property T

Applications

Theorem *The arrangement of n subspaces \mathcal{H}_i satisfying the conditions*

$$\angle(\mathcal{H}_i, \mathcal{H}_j) \geq \alpha_{ij}$$

for all i, j , is “good” if there exists an n -dimensional spherical simplex with angles between the faces equal to α_{ij} .

Actually, “if” can be replaced with “if and only if”, but the other direction is much easier.

“Proof”

Expander Graphs

Kazhdan Constants

Computation of Kazhdan Constants

Bounded Generation Property

Kazhdan Constants for $SL_n(\mathbb{F}_p)$

Using Geometry to (dis-)Prove Property T

Subspace Arrangements

Easy Example

- An Example
- Main result
- Equivalent Statement
- “Proof”
- Application to property T

Applications

It seems “obvious” that the most difficult case is when \mathcal{H}_i are hyper-planes (they have co-dimension 1 in \mathcal{H}).

In this case WLOG we can assume that $\dim \mathcal{H} = n$ and we can find unit vectors u_i perpendicular to the hyper-planes \mathcal{H}_i . By construction we have $|\langle u_i, u_j \rangle| \leq \varepsilon_{i,j}$.

The volume of the simplex spanned by u_i is

$$V^2 = \det(\langle u_i, u_j \rangle)_{i,j} \geq \det A$$

is bounded from below, thus any interior point is “far” from at least one face, which translates to a lower bound for the angle between the subspaces \mathcal{H}_i .

In the general case one need to do a clever induction to justify the “obvious”.

Application to property T

Expander Graphs

Kazhdan Constants

Computation of Kazhdan Constants

Bounded Generation Property

Kazhdan Constants for $SL_n(\mathbb{F}_p)$

Using Geometry to (dis-)Prove Property T

Subspace Arrangements

Easy Example

- An Example
- Main result
- Equivalent Statement
- “Proof”
- Application to property T

Applications

Theorem (Dymara-Januszkiewicz) *Let G be a group generated by a finite subgroup G_i . If for any i, j the group $G_{i,j} = \langle G_i, G_j \rangle$ has property T and the Kazhdan constant $\mathcal{K}(G_{ij}, G_i \cup G_j)$ is sufficiently big, then G has property T.*

Expander Graphs

Kazhdan Constants

Computation of Kazhdan Constants

Bounded Generation Property

Kazhdan Constants for $SL_n(\mathbb{F}_p)$

Using Geometry to (dis-)Prove Property T

Subspace Arrangements

Easy Example

Applications

- Coxeter Groups
- $SL_n(\mathbb{F}_p[t_1, t_2])$
- Groups graded by root systems

Applications

Coxeter Groups

Expander Graphs

Kazhdan Constants

Computation of Kazhdan
Constants

Bounded Generation
Property

Kazhdan Constants for
 $SL_n(\mathbb{F}_p)$

Using Geometry to
(dis-)Prove Property T

Subspace Arrangements

Easy Example

Applications

- **Coxeter Groups**
- $SL_n(\mathbb{F}_p[t_1, t_2])$
- Groups graded by root systems

Theorem (K.) *A Coxeter group G has property T if and only if it is finite. Moreover, the Kazhdan constant $\mathcal{K}(G, S)$ can be computed using the defining representation of the group G .*

Corollary *The mixing time of the random walk on a Coxeter group G with respect to the standard generating set S is bounded by $n^3 \log n$, where n is the rank of G .*

$SL_n(\mathbb{F}_p[t_1, \dots, t_k])$

Expander Graphs

Kazhdan Constants

Computation of Kazhdan Constants

Bounded Generation Property

Kazhdan Constants for $SL_n(\mathbb{F}_p)$

Using Geometry to (dis-)Prove Property T

Subspace Arrangements

Easy Example

Applications

- Coxeter Groups
- $SL_n(\mathbb{F}_p[t_1, t_2])$
- Groups graded by root systems

Theorem *The groups $SL_n(\mathbb{F}_p[t_1, \dots, t_k])$ have property T for any $n \geq 3$ and $p > 4$.*

Proof: The subgroups

$$G_i = \text{Id} + \mathbb{F}_p E_{i,i+1}$$

$$\begin{pmatrix} 1 & G_1 & 0 & 0 \\ 0 & 1 & G_2 & 0 \\ 0 & 0 & 1 & G_3 \\ G_4 & 0 & 0 & 1 \end{pmatrix}$$

and

$$G_n = \text{Id} + (\mathbb{F}_p + t_1 \mathbb{F}_p + \dots + t_k \mathbb{F}_p) E_{n,1}$$

generate the group $SL_n(\mathbb{F}_p[t_1, \dots, t_k])$.

Any two of these group either commute or generate a Heisenberg group mod p .

Expander Graphs

Kazhdan Constants

Computation of Kazhdan Constants

Bounded Generation Property

Kazhdan Constants for $SL_n(\mathbb{F}_p)$

Using Geometry to (dis-)Prove Property T

Subspace Arrangements

Easy Example

Applications

- Coxeter Groups
- $SL_n(\mathbb{F}_p[t_1, t_2])$
- Groups graded by root systems

Thus the "matrix" of cosines of the angles between the subspaces of fixed points is:

$$\begin{pmatrix} 1 & -\varepsilon & 0 & \dots & -\varepsilon \\ -\varepsilon & 1 & -\varepsilon & \dots & 0 \\ 0 & -\varepsilon & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ -\varepsilon & 0 & 0 & \dots & 1 \end{pmatrix}$$

where $\varepsilon = p^{-1/2}$.

This matrix is positive definite if $\varepsilon < 1/2$.

Same result holds for Steinberg groups over any finitely generated ring of characteristic $p \geq 10$.

Groups graded by root systems

Expander Graphs

Kazhdan Constants

Computation of Kazhdan Constants

Bounded Generation Property

Kazhdan Constants for $SL_n(\mathbb{F}_p)$

Using Geometry to (dis-)Prove Property T

Subspace Arrangements

Easy Example

Applications

- Coxeter Groups
- $SL_n(\mathbb{F}_p[t_1, t_2])$
- Groups graded by root systems

Definition Let Φ be a finite root system. A group G is called graded by Φ if there exists root subgroup X_α for $\alpha \in \Phi$ such that

- G is generated by X_α
- if α and β are linearly independent then $[X_\alpha, X_\beta] \subset \langle X_\gamma \rangle$, where $\gamma = a\alpha + b\beta$ for $a, b \geq 1$
- if $\beta = a\alpha$ for $a \geq 1$ then $[X_\alpha, X_\beta] \subset \langle X_\gamma \rangle$, where $\gamma = b\beta$ for $b \geq 1$
- for each $\alpha \in \Phi$ there exists a set of positive roots Φ^+ such that $\alpha \in \Phi^+$ and

$$X_\alpha \subset \langle X_\gamma \mid \gamma \in \Phi^+, \gamma \neq \alpha \rangle.$$

Expander Graphs

Kazhdan Constants

Computation of Kazhdan Constants

Bounded Generation Property

Kazhdan Constants for $SL_n(\mathbb{F}_p)$

Using Geometry to (dis-)Prove Property T

Subspace Arrangements

Easy Example

Applications

- Coxeter Groups
- $SL_n(\mathbb{F}_p[t_1, t_2])$
- Groups graded by root systems

Theorem (Ershov-Jaikin-K.) *Let G be a group graded by the root system Φ . Then the Kazhdan constant $\mathcal{K}(G, \cup X_\alpha) > 0$, i.e., G almost has property T.*

The proof is a bit messy...

Corollary *Let $G = St_\Phi(R)$ be a (twisted) Steinberg group of rank ≥ 2 over a finitely generated ring R .*

Then G has property T and there is an lower bound for the Kazhdan constant $\mathcal{K}(G, S) > 0$, where S is the "natural" generating set of G .